arXiv:1112.4137v1 [math-ph] 18 Dec 2011

Sigma functions for a space curve $(3, 4, 5)$ type with an appendix by J. Komeda

SHIGEKI MATSUTANI

Abstract. In this article, a generalized Kleinian sigma function for an affine $(3, 4, 5)$ space curve of genus 2 was constructed as the simplest example of sigma function for an affine space curves, by applying the construction method of sigma function for (r, s) plane curve provided by Eilbeck Enolskii and Leykin (SIDEIII, CRM Proc. Lecture Notes, 25 2000) to the space curve. By defining the fundamental differential of the second kind over it, the Legendre relations was obtained as the symplectic structure over it. It was showed that with the abelian map to $\mathbb{C}^2$, the symplectic structure determines the sigma function. Using the sigma function, the Jacobi inversion formulae for the curve are obtained. It means that the generalization of the sigma functions for the affine plane curves to ones for the space curves is basically possible. An interesting relation between a semigroup generated by $(6, 13, 14, 15, 16)$ and Norton number associated with Monster group is also mentioned with an Appendix by Komeda.

sigma function, space curve, Jacobi inversion formula

MSC 2010: 14H05, 14H42, 14H50, 14H55, 20M07

1. Introduction

In Tata lectures on theta II [Mu], Mumford polished the abelian function theory in the nineteenth century on the hyperelliptic curves based upon the study of Jacobi, from a viewpoint of modern algebraic geometry, and investigated the abelian functions of hyperelliptic curves. Using the explicit expressions, he showed that the abelian functions are closely connected to the nonlinear integrable systems, kinematic problems, representation theory, and so on. He described these explicit connections in terms of the theta functions.

Recently on hyperelliptic curves, Grand [Gr], Ônishi [O], and Buchstaber, Enolskii, and Leykin [BEL1] found that it is more convenient to show the abelian functions in terms of the Kleinian sigma function which is a natural generalization of the Weierstrass sigma function [B1, B2, K, W] and investigated the hyperelliptic functions using it. For example, in terms of the sigma functions, it is much easier to rewrite formulae in [Mu] than theta function [BEL1, Ma01].

Further in [EEL], Enolskii, Eilbeck, and Leykin discovered a construction which generalizes the Kleinian sigma function for a (r, s) plane curve of genus g, though the original

---

Date:

Kleinian sigma function is only for a hyperelliptic curve; r and s ($r < s$) are coprime positive integers $g = (r-1)(s-1)/2$. In [EEL], they, firstly, constructed the fundamental differential of the second kind over an affine (r, s) plane curve and using it, obtained the Legendre relations as the symplectic structure over the curve. Using the Legendre relations, they defined the generalized Kleinian sigma function over $\mathbb{C}^g$, the image of the abelian map. They found the natural Jacobi inversion formulae in terms of their sigma function. We call the construction *EEL construction* in this article. Using the EEL construction, Buchstaber, Enolskii, Leykin, Eilbeck, Ônishi, Nakayashiki, Previato, and Matsutani also have studied abelian functions over (r, s) affine curves [BEL2, BLE1, BLE2, EEMOP1, EEMOP2, N, Ma01, MP08, MP09, MP11]. One of these purposes is also to generalize Mumford's investigations in [Mu] and make them more profound. These results are connected with many physical problems e.g., [BG, BDE, MP09].

In this article, we consider a generalized Kleinian sigma function for an affine $(3, 4, 5)$ space curve of genus 2, which is the simplest affine space curve. Our purpose of this article is to show that sigma function is also defined for an affine space curve as we can do for plane curves.

Following the EEL-construction, we define the fundamental differential of the second kind over it and obtain the Legendre relations as the symplectic structure over it. With the abelian map to $\mathbb{C}^2$, we show that the symplectic structure determines the sigma function. Further using the sigma function, we obtain the Jacobi inversion formulae for the curve and the Jacobian following the previous work [MP08].

It means that the generalization of the sigma functions for the affine plane curves to ones for the space curves is basically possible and is useful.

When a physical phenomenon is described by a (r, s) plane curve including $r = 2$ case, we naturally encounter a singular (r, s) plane curve because the phenomenon is governed by the parameter space of the curve and the variation of the parameters includes ones for singular curves. The singular curves sometimes become space curves by normalization as in Example 3 in §2.2. Hence we believe that this demonstration that we have sigma function for a space curve is quite crucial.

In Discussion we also show a problem of a space curve associated with the semigroup generated by $(6, 13, 14, 15, 16)$ with an Appendix by Komeda. The semigroup might be related to Norton number associated with the Monster group, the simple largest sporadic finite group [Mc, MS, N]. In fact, the gap sequence of the semigroup is $\{1, 2, 3, 4, 5, 7, 8, 9, 10, 11, 17, 23\}$ whereas Norton number is $\{1, 2, 3, 4, 5, 7, 8, 9, 11, 17, 19, 23\}$ [Cu, Mc, MS, N]. Both sequences agree except 19 and 10. The gap sequence is related to the Sato weight of the KP hierarchy whereas Norton number is the weight of the replicable functions which govern the moonshine phenomena of the Monster group. The sigma function for the space curve is similar to the replicable function [KMP]. We give some comments on these relations.

I thank John McKay for posing my attention to the Norton problem and his encouragement and Jiryo Komeda for crucial discussions on semigroup and for his Appendix on

the semigroup $\langle 6, 13, 14, 15, 16\rangle$. This work started in a seminar at Yokohama National university 2008, and was stimulated by the international conference at HWK 2011. I am grateful to Kenichi Tamano, Norio Konno, Claus Lämmerzahl, Jutta Kunz, and Victor Enolskii. I am also most grateful to Emma Previato for crucial discussions on the preparation [KMP].

2. Mathematical Preliminary

2.1. **Numerical semigroup.** In this section, we give an overview of recent study of the numerical semigroups as sub-semigroup of non-negative integers $\mathbb{N}_0$. We call an additive semigroup in $\mathbb{N}_0$ *numerical semigroup* if its complement in $\mathbb{N}_0$ is finite.

For a numerical semigroup $H = H(M)$ generated by M, the number of elements of $L(H) := \mathbb{N}_0 \setminus H$ is called genus. For example, we have semigroups H_2, H_4, H_{12} generated by

$$M_2 := \langle 3, 4, 5\rangle, \quad M_4 := \langle 3, 7, 8\rangle, \quad M_{12} := \langle 6, 13, 14, 15, 16\rangle,$$

respectively whose genera are $g(H_g)$ for $g = 2, 4, 12$ due to

$$L(H_2) := \{1, 2\}, \quad L(H_4) := \{1, 2, 4, 5\}, \quad L(H_{12}) := \{1, 2, 3, 4, 5, 7, 8, 9, 10, 11, 17, 23\}.$$

For a gap sequence $L := \{\ell_0 < \ell_1 < \cdots < \ell_{g-1}\}$ of genus g, let $M(L)$ be the minimal set of generators for the semigroup $H(L)$ and

$$a(L) := \{a_0(L), a_1(L), \cdots, a_{g-1}(L)\}$$

where $a_i := \ell_i - i - 1$. Hence we have

$$a(L(H_2)) = \{0^2\}, \quad a(L(H_4)) = \{0^2, 1^2\} \quad \text{and} \quad a(L(H_{12})) := \{0^5, 1^5, 6, 11\}.$$

Let $a_{\min}(L)$ be the smallest positive integer of $M(L)$, and $wt(L) := \sum_{i=0}^{g-1} a_i(L)$ be the weight of L; Since $a_{\min}(L)$ plays an essential role in semigroup, we call semigroup with $a_{\min}(L)$ "$a_{\min}(L)$-semigroup"; H_2 and H_4 are 3-semigroups and H_{12} is a 6-semigroup.

For a complete non-singular irreducible curve C of genus g over an algebraically closed field k of characteristic 0, the field of its rational functions $k(C)$, and a point $P \in C$, we define

$$H(P) := \{n \in \mathbb{N}_0 \mid \text{there exists } f \in k(C) \text{ such that } (f)_\infty = nP \}, \tag{2.1}$$

which is called the Weierstrass semigroup of the point P. If $L(H(P)) := \mathbb{N}_0 \setminus H(P)$ differs from the set $\{1, 2, \cdots, g\}$, we call P Weierstrass point of C.

A numerical semigroup H is said to be Weierstrass if there exists a pointed algebraic curve (C, P) such that $H = H(P)$. Hurwitz considered whether every numerical semigroup H is Weierstrass. This was a long-standing problem but Buchweitz finally showed that every H is not Weierstrass. His first counterexample is the semigroup H_B generated by 13, 14, 15, 16, 17, 18, 20, 22 and 23, whose genus is 16.

Thus in general, it is not so trivial whether a given semigroup is Weierstrass or not. Komeda has been investigated this problem with Ohbuchi [K83, K99, K04, KO04, KO08, KO08a]. Komeda and his coauthors, Hensel, Landsberg, Schaps, Pinkham, Maclahlan, Eisenbud and Harris showed that the following is Weierstrass;

(1) semigroup whose cardinality of the generators is less than 4,
(2) semigroup whose genus is less than 9,
(3) semigroup whose all primitive sequence, twice the smallest positive integer in $H(L) >$ the largest integer in L, of genus 9,
(4) semigroup whose $a_{\min}(L) = 2, 3, 4, 5$,
(5) semigroup whose $a(L) = (0^{g-2}, m, n)$ for genus g and $wt(L) = g$,
(6) semigroup whose $a(L) = (0^{g-r}, m^r)$ for genus g, and
(7) other many special cases of n-semigroups generated by 4 elements.

2.2. **Commutative Algebra.** We review normal ring and normalization in commutative ring [Mat]. We assume that every ring is a commutative ring with unit.

B is a ring and A is a subring of B. B is said to be *extension* of A. An element b of B is said to be *integral* over A if b satisfies a monic polynomial over A, i.e., there exist n and $\{a_i\}_{i=1,\dots,n} \in A$ such that

$$b^n + a_1 b^{n-1} + \cdots a_n = 0.$$

We say that B is *integral* over A, or B is an *integral ring* over A, or B is an *integral extension* of A if every element b of B is integral over A.

An *integral closure* in B over A is defined by

$$\tilde{A} := \{b \in B \mid a \text{ is integral over } A\}.$$

If $A = \tilde{A}$, A is *integral closed* in B.

Definition 2.1. *A is a ring and $Q(A)$ is a quotient ring of A. We assume that A is an integral domain. A is* normal *if A is integral closed in $Q(A)$, i.e., for $\tilde{A} := \{q \in Q(A) \mid$ there exist n and $a_i \in A$ such that $q^n + a_1 q^{n-1} + \cdots a_n = 0$ $\}$, $A = \tilde{A}$.*

We define the minimum extension $\hat{A}$ of A in $Q(A)$ so that $\hat{A}$ is integral closed in $Q(A)$. We say that $\hat{A}$ is normalization *of A or the* normalized ring *of A.*

Example 1 : $x^3 - y^2$: $R := \mathbb{C}[X, Y]/(X^3 - Y^2)$ is not normal because $\frac{Y}{X} \in \tilde{R} \setminus R \subset Q(R)$ due to $\left(\frac{Y}{X}\right)^2 - X = 0$. Since $R \approx \mathbb{C}[t^2, t^3]$, the normalized ring is $\hat{R} = \mathbb{C}[t]$.

Example 2 : $y^3 = x^5 - 1$ and $w^3 = z - z^6$: Following the statements in [ACGH, p.31], we consider the covering of a curve of $f(x, y) := y^3 - x^5 + 1$. Let us consider a homogeneous polynomial $F(X, Y, Z) := Y^3 Z^2 - X^5 + Z^5 \in \mathbb{C}[X, Y, Z]$. Around $Z \neq 0$, we have

$$F(X, Y, Z) = Z^5 \left(\frac{Y^3}{Z^3} - \frac{X^5}{Z^5} + 1\right)$$

and thus by regarding that $x = X/Z$ and $y = Y/Z$, we have $F(X, Y, Z) = Z^5 f(X/Z, Y/Z)$. $R_0 := \mathbb{C}[x, y]/(f(x, y))$ is a normal ring. On the other hand, around $Z = 0$ and $X \neq 0$, we have

$$F(X,Y,Z) = X^5\left(\frac{Y^3Z^2}{X^5} - 1 + \frac{Z^5}{X^5}\right),$$

and then we obtain a polynomial, $g(w,z) = w^3z^2 - 1 + z^5$ by regarding $w = Y/X$ and $z = Z/X$. However $R_\infty := \mathbb{C}[w,z]/(g(w,z))$ is not a normal ring. As a vector space, R_∞ is

$$\begin{aligned} R_\infty = &\mathbb{C}1 + \mathbb{C}z + \mathbb{C}z^2 + \mathbb{C}z^3 + \mathbb{C}z^4 + \cdots \\ &+ \mathbb{C}w + \mathbb{C}wz + \mathbb{C}wz^2 + \mathbb{C}wz^3 + \mathbb{C}wz^4 + \cdots \\ &+ \mathbb{C}w^2 + \mathbb{C}w^2z + \mathbb{C}w^2z^2 + \mathbb{C}w^2z^3 + \mathbb{C}w^2z^4 + \cdots \\ &+ \mathbb{C}w^3 + \mathbb{C}w^3z. \end{aligned}$$

We will show that $q \in Q(R_\infty) \setminus R_\infty$ exists such that $q^n + a_1q^{n-1} + \cdots a_n = 0$ for certain $a_i \in R_\infty$. Noting

$$\frac{1}{1-z}g(w,z) = \frac{w^3z^2}{1-z} + 1 + z + z^2 + z^3 + z^4 = 0 \in Q(R_\infty),$$

we consider

$$q := \frac{w^3}{1-z} + \frac{1+z}{z^2} \in Q(R_\infty) \setminus R_\infty,$$

which is integral over R_∞. By normalization, we define $\hat{w} := wz = y/x^2$. $\hat{R}_\infty := \mathbb{C}[\hat{w},z]/(\hat{g}(w,z))$ is a normal ring, where

$$\hat{g}(\hat{w},z) := \hat{w}^3 - z + z^6.$$

The minimal condition is obvious.

Example 3 : a space curve; $y^3 = x^2(x^2-1)$ and $w^3 = x(x^2-1)^2$: Let us consider a polynomial $f(x,y) = y^3 - x^2(x^2-1)$ and show that $R_0 := \mathbb{C}[x,y]/(f(x,y))$ is not a normal ring. As a vector space, R_0 is

$$\begin{aligned} R_0 = &\mathbb{C}1 + \mathbb{C}x + \mathbb{C}x^2 + \mathbb{C}x^3 + \mathbb{C}x^4 + \cdots \\ &+ \mathbb{C}y + \mathbb{C}yx + \mathbb{C}yx^2 + \mathbb{C}yx^3 + \mathbb{C}yx^4 + \cdots \\ &+ \mathbb{C}y^2 + \mathbb{C}y^2x + \mathbb{C}y^2x^2 + \mathbb{C}y^2x^3 + \mathbb{C}y^2x^4 + \cdots. \end{aligned}$$

We show that $w \in Q(R_0) \setminus R_0$ exists such that $w^n + a_1w^{n-1} + \cdots a_n = 0$ for certain a_i's of R_0. In other words, noting

$$y \sim \sqrt[3]{x^2(x^2-1)} \quad \text{and} \quad y^2 \sim x\sqrt[3]{x(x^2-1)},$$

one of w is that $w := \frac{y^2}{x}$ which is integral over R_0 because $w^3 = \frac{y^6}{x^3} = x(x^2-1)^2$ or $w^3 - x(x^2-1)^2 = 0 \in R_0$. Let $g(x,w) = x(x^2-1)^2$. Noting the relations,

$$w = \frac{y^2}{x}, \quad y = \frac{w^2}{x^2-1}, \quad wy = (x^2-1)x,$$

we have the normalized ring of R_0,

$$\hat{R}_0 := \mathbb{C}[x, y, w]/(f_1(x, y, z), f_2(x, y, z)f_3(x, y, z)),$$

where

$$f_1(x, y, w) = y^2 - wx, \quad f_2(x, y, w) = wy - (x^2 - 1)x, \quad \text{and} \quad f_3(x, y, w) = w^2 - y(x^2 - 1).$$

The minimal condition is also obvious. This Example 3 corresponds to the special case of the affine $(3, 4, 5)$ space curve in this article.

3. A CURVE (3,4,5)

Since H_2 generated by $\langle 3, 4, 5\rangle$ is Weierstrass and is the simplest semigroup whose cardinality of the generators is greater than 2, we consider a curve $C(H_2)$ explicitly in order to construct the sigma functions for $C(H_2)$ following the EEL construction.

In order to construct a non-singular curve $X_2 = C(H_2)$, we consider two singular curves X_3 and X_4 generated by the zeroes of

$$f_{3,12}(x, y_4) := y_4^3 - k_4(x), \quad k_4(x) := k_2(x)k_1(x)^2,$$

and

$$f_{4,15}(x, y_5) := y_5^3 - k_5(x), \quad k_5(x) := k_2(x)^2 k_1(x),$$

where for finite $b_a \in \mathbb{C}$ $(a = 1, 2, 3)$ which is distinct from each other, and

$$\begin{aligned} k_2(x) &:= (x - b_1)(x - b_2) = x^2 + \lambda_1^{(2)} x + \lambda_2^{(2)}, \\ k_1(x) &:= (x - b_0) = x + \lambda_1^{(1)}. \end{aligned}$$

Let us consider commutative rings $R_3 := \mathbb{C}[x, y_4]/(f_{3,12}(x, y_4))$ and $R_4 := \mathbb{C}[x, y_5]\ /(f_{4,15}(x, y_5))$. These arithmetic genera are three and four respectively, though the geometric genera are not. Following the normalization in subsection 2.2, we normalize R_3 and R_4. Since in terms of the language of the commutative algebra [Mat], $\dfrac{y_4^2}{(x - b_0)}$ is integral over R_3 in $Q(R_3)$ and $\dfrac{y_5^2}{(x - b_0)(x - b_2)}$ is integral over R_4 in $Q(R_4)$, R_3 and R_4 are not normal rings. Thus we will normalise them in $\mathbb{C}[x, y_4, y_5]$ in the meaning of the commutative algebra [Mat] (see Example 3 in §2.2).

For the zeroes of $f_{3,12}(x, y_4)$ and $f_{3,15}(x, y_4)$, we could have the relations,

$$y_4 y_5 = k_2(x)k_1(x), \quad y_5 = \frac{y_4^2}{(x - b_0)}, \quad y_4 = \frac{y_5^2}{(x - b_1)(x - b_2)}. \tag{3.1}$$

Here for the primitive root ζ_3 ($\zeta_3^3 = 1, \zeta_3 \neq 1$), ζ_3 acts on X_3 and X_4 respectively. The first relation is chosen in the possibilities $y_4 y_5 = \zeta_3^i k_2(x)k_1(x)$ $i = 0, 1, 2$.

As a normalisation of these singular curves, we consider the commutative ring,

$$R_2 \equiv R := \mathbb{C}[x, y_4, y_5]/(f_8, f_9, f_{10}),$$

and $X_2 := \text{Spec } R$. Here we define $f_8, f_9, f_{10} \in \mathbb{C}[x, y_4, y_5]$ by

$$f_8 = y_4^2 - y_5 k_1(x), \quad f_9 = y_4 y_5 - k_2(x) k_1(x), \quad f_{10} = y_5^2 - y_4 k_2(x),$$

which are also regarded as the 2×2 minors of

$$\left| \begin{matrix} k_2(x) & y_4 & y_5 \\ y_4 & y_5 & k_3(x) \end{matrix} \right|. \tag{3.2}$$

Here ζ_3 acts on X_2 by $\hat{\zeta}_3(x, y_4, y_5) = (x, \zeta_3 y_4, \zeta_3^2 y_5)$.

Let X be the Riemann surface which is naturally obtained as an extension of X_2 as mentioned in [ACGH, p.31], i.e., $X = X_2 \cup \{\infty\}$ as a set. It is noted that when x diverges, y_4 and y_5 also diverge vise versa. Thus the infinity point ∞ uniquely exists in X. $\mathbb{G}_m$ acts on R by setting $g_m^{-3} x$, $g_m^{-a} y_a$ for x, y_a, $g_m \in \mathbb{G}_m$ and $a = 4, 5$. By Nagata's Jacobi-method [Mat], it can be proved that X is non-singular.

Though they do not explicitly appear, we will also implicitly consider parametrisations of y_4 and y_5 by

$$y_4 = w_2 w_1^2, \quad y_5 = w_2^2 w_1, \tag{3.3}$$

where

$$w_1^3 = k_1, \quad w_2^3 = k_2. \tag{3.4}$$

When we consider $\tilde{R} := \mathbb{C}[x, w_1, w_2]/(w_1^3 - k_1(x), w_2^3 - k_2(x))$, it is related to the principal ideal theorem for X and a natural covering of X.

3.1. **The Weierstrass gap and holomorphic one forms.**

The Weierstrass gap sequences at ∞ are given by the following Table 1. For the local parameter t_∞ at ∞, we have

$$x = \frac{1}{t_\infty^3}, \quad y_4 = \frac{1}{t_\infty^4}(1 + d_{\geq}(t_\infty)), \quad y_5 = \frac{1}{t_\infty^5}(1 + d_{\geq}(t_\infty)), \tag{3.5}$$

Here for a given local parameter t at some P in X, by $d_{\geq}(t^\ell)$ we denote the terms of a function on X in its t-expansion whose orders of zero at P are greater than ℓ or equal to ℓ. $H(\infty)$ in (2.1) is $H(3, 4, 5)$.

Pinkham considered $(3, 4, 5)$ curve as the simplest example of numerical semigroup [P, Sec.14]. Its monomial curve is defined by,

$$Z_4^2 = Z_3 Z_5, \quad Z_4 Z_5 = Z_3^5, \quad Z_5^2 = Z_3^3 Z_4, \quad \text{or the } 2 \times 2 \text{ minor of } \left| \begin{matrix} Z_3 & Z_4 & Z_5 \\ Z_4 & Z_5 & Z_3^2 \end{matrix} \right|.$$

Z_3, Z_4 and Z_5 correspond to $\dfrac{1}{x}$ $\dfrac{1}{y_4}$ and $\dfrac{1}{y_5}$ respectively and these relations correspond to (3.1) and (3.2).

Table 1

	0	1	2	3	4	5	6	7	8	9	10	11	12	13	8	9	10
X_3	1	-	-	x	y_4	-	x^2	xy_4	y_4^2	x^3	x^2y_4	xy_4^2	x^4	x^3y_4	$x^2y_4^2$	x^5	x^4y_4
X_4	1	-	-	x	-	y_5	x^2	-	xy_5	x^3	y_5^2	x^2y_5	x^4	x^2y_5	xy_5^2	x^5	$x^2y_5^2$
X_2	1	-	-	x	y_4	y_5	x^2	xy_4	xy_5	y_4y_5	x^2y_4	x^2y_5	x^4	x^3y_4	x^3y_5	x^5	x^4y_4

There we define $\phi_i^{(g)}$ as a non-gap monomial in R_g for $g = 2, 3, 4$ and *e.g.*, $\phi_0^{(2)} = 1$, $\phi_1^{(2)} = x$, $\phi_2^{(2)} = y_4$, $\phi_3^{(2)} = y_5$, $\phi_4^{(2)} = x^2$, $\cdots$ and $\phi_0^{(3)} = 1$, $\phi_1^{(3)} = x$, $\phi_2^{(3)} = y_4$, $\phi_3^{(3)} = x^2$, $\phi_4^{(3)} = xy_4$, $\cdots$. We introduce the weight $N^{(g)}(n)$ by letting $N^{(g)}(n) := -\mathrm{wt}(\phi_n^{(g)})$, where wt() is the degree of divisor at ∞ of each curve X's. It is noted that H_2 is identical to $\{N^{(2)}(n) \mid n = 0, 1, 2, \ldots\}$. By letting

$$\Lambda_i^{(g)} := N^{(g)}(g) - N^{(g)}(i-1) - g + i - 1,$$

the related Young diagrams $\Lambda \equiv \Lambda^{(2)} := (\Lambda_1, \Lambda_2) = (1, 1)$, $\Lambda^{(3)} := (\Lambda_1^{(3)}, \Lambda_2^{(3)}, \Lambda_3^{(3)}) = (3, 1, 1)$ and $\Lambda^{(4)} := (\Lambda_1^{(4)}, \Lambda_2^{(4)}, \Lambda_3^{(4)}, \Lambda_4^{(4)}) = (4, 2, 1, 1)$ are given by respectively:



The Young diagram Λ is not symmetric, whereas ${}^t\Lambda^{(3)} = \Lambda^{(3)}$ and ${}^t\Lambda^{(4)} = \Lambda^{(4)}$.

Here we note that the Young diagram is related to the quantities in the semigroup as

$$\Lambda_{g-i}^{(g)} = a_i(L(\tilde{H}_g)) + 1, \quad |\Lambda^{(g)}| = \sum_{i=1}^{g} \Lambda_i^{(g)} = wt(L(\tilde{H}_g))) + g,$$

where $\tilde{H}_2 = H_2$, $\tilde{H}_3 = H(3, 4)$ and $\tilde{H}_4 = H(3, 5)$.

Proposition 3.1. *Bases of the holomorphic one forms over X can be expressed by*

$$\nu_1^I := \frac{dx}{3y_5}, \quad \nu_2^I := \frac{dx}{3y_4}.$$

For later convention, we introduce $\phi_{H^1 i} \in R$ $(i = 1, 2, 3, \cdots)$ by

$$\phi_{H^1 0} := y_4, \quad \phi_{H^1 1} := y_5, \quad \phi_{H^1 2} := xy_4, \quad \phi_{H^1 3} := xy_5,$$

for $i > 3$,

$$\phi_{H^1 i} := \begin{cases} x^{(i-4)/3} y_4 y_5 & i \equiv 1 \bmod 3, \\ x^{(i+1)/3} y_4 & i \equiv 2 \bmod 3, \\ x^{i/3} y_5 & i \equiv 0 \bmod 3. \end{cases}$$

Then we have

$$\nu_i^I := \frac{\phi_{{H^1}_{i-1}} dx}{3 y_4 y_5}, \quad (i = 1, 2).$$

We also define the weight $N_{H^1}(n)$ by $N_{H^1}(n) := -\mathrm{wt}(\phi_{{H^1}_n})$; $N_{H^1}(0) = 4$, $N_{H^1}(1) = 5$, and $N_{H^1}(n) = n + 5$ for $n \geq 2$.

Then the following proposition is obvious.

Proposition 3.2. *$\sum_{i=0}^n a_i \tilde{\nu}_i$ belongs to $H^1(X \setminus \infty, \mathcal{O}_X)$, where $\tilde{\nu}_i := \frac{\phi_{{H^1}_i} dx}{3 y_4 y_5}$ and the order of the singularity of $(\tilde{\nu}_i)$ at ∞ is given by $N_{H^1}(n) - 5$.*

Lemma 3.3.

$$a_0 \frac{dx}{y_4 y_5} + a_1 \frac{x dx}{y_4 y_5} + a_2 \frac{x^2 dx}{y_4 y_5}$$

is not holomorphic one form over X if a_i does not vanish.

Proof. For $n < 3$, every $\sum_{i=0}^n a_i \frac{x^i dx}{y_4 y_5}$ has singularities at points in $X \setminus \infty$. □

We choose the bases α_i, β_j $(1 \leqq i, j \leqq 2)$ of $H_1(X, \mathbb{Z})$ such that their intersection numbers are $\alpha_i \cdot \alpha_j = \beta_i \cdot \beta_j = 0$ and $\alpha_i \cdot \beta_j = \delta_{ij}$, and we denote the period matrices by

$$[\omega' \ \omega''] = \frac{1}{2} \left[\int_{\alpha_i} \nu_j^I \quad \int_{\beta_i} \nu_j^I \right]_{i,j=1,2}. \tag{3.6}$$

Let Π_2 be a lattice generated by ω' and ω''. For a point $P \in X$, the abelian map $w : X \to \mathbb{C}^2$ is defined by

$$w(P) = \int_\infty^P \nu^I \in \mathbb{C}^2,$$

and for a point $(P_1, \cdots, P_k) \in S^k X$, i.e., the k-th symmetric product of X, the abelian map $w : S^k X \to \mathbb{C}^2$ by $w(P_1, \cdots, P_k) := \sum_{i=1}^k w(P_i)$. Then we define the Jacobian $\mathcal{J}_2$ and its subvariety $\mathcal{W}_k$ $(k = 0, 1, 2)$ by

$$\kappa : \mathbb{C}^2 \to \mathcal{J}_2 = \mathbb{C}^2 / \Pi_2 = \mathcal{W}_2, \quad \mathcal{W}_k := \kappa w(S^k X)$$

respectively.

For a point $(P_1, P_2) \in S^2 X$ around the infinity point, by letting their local parameters $t_{\infty,1}$ and $t_{\infty,2}$ $u \equiv {}^t(u_1, u_2) := w(P_1, P_2)$ is given by

$$\begin{aligned} u_1 &= \frac{1}{2}(t_{\infty,1}^2 + t_{\infty,2}^2)(1 + d_{>0}(t_{\infty,1}, t_{\infty,2})), \\ u_2 &= (t_{\infty,1} + t_{\infty,2})(1 + d_{>0}(t_{\infty,1}, t_{\infty,2})), \end{aligned} \tag{3.7}$$

where $d_{\geq}(t_1, t_2)$ is a natural extension of $d_{\geq}(t)$.

3.2. **Differentials of the second and the third kinds.** Following the EEL-construction [EEL, BEL2] for a (n, s) curve, we give an algebraic representation of a differential form which, up to a tensor of holomorphic one forms, is equal to the fundamental normalized differential of the second kind in [F, Corollary 2.6]. In other words, we apply the EEL-construction to the space curve X.

Definition 3.4. *A two-form $\Omega(P_1, P_2)$ on $X \times X$ is called a fundamental differential of the second kind if it is symmetric,*

$$\Omega(P_1, P_2) = \Omega(P_2, P_1), \tag{3.8}$$

it has its only pole (of second order) along the diagonal of $X \times X$, and in the vicinity of each point (P_1, P_2) is expanded in power series as

$$\Omega(P_1, P_2) = \Big(\frac{1}{(t_{P_1} - t'_{P_2})^2} + d_{\ge}(1)\Big) dt_{P_1} \otimes dt_{P_2} \quad \text{(as } P_1 \to P_2\text{)}, \tag{3.9}$$

where t_P is a local coordinate at the point $P \in X$.

Here we use the convention that for $P_a \in X$, P_a is represented by $(x_a, y_{4,a}, y_{5,a})$ or $(x_{P_a}, y_{4,P_a}, y_{5,P_a})$ and for $P \in X$, P is expressed by (x, y_4, y_5).

Proposition 3.5. *Let $\Sigma\big(P, Q\big)$ be the following form,*

$$\Sigma\big(P, Q\big) := \frac{y_{4,P}y_{5,P} + y_{4,P}y_{5,Q} + y_{4,Q}y_{5,P}}{(x_P - x_Q)3y_{4,P}y_{5,P}} dx_P. \tag{3.10}$$

Then $\Sigma(P, Q)$ has the properties

1. *$\Sigma(P, Q)$ as a function of P is singular at $Q = (x_Q, y_{4,Q}, y_{5,Q})$ and ∞, and vanishes at $\hat{\zeta}_3^{\ell}(Q) = (x_Q, \zeta_3^{\ell} y_{4,Q}, \zeta_3^{2\ell} y_{5,Q})$, $(\ell = 1, 2)$.*
2. *$\Sigma(P, Q)$ as a function of Q is singular at P and at ∞.*

Proof. Direct computations lead the results. □

The following holds for non-singular X;

Proposition 3.6. *There exist differentials $\nu_j^{II} = \nu_j^{II}(x, y_4, y_5)$ $(j = 1, 2)$ of the second kind such that they have their only pole at ∞ and satisfy the relation,*

$$\begin{aligned} &d_Q\Sigma\big(P, Q\big) - d_P\Sigma\big(Q, P\big) \\ &\qquad = \sum_{i=1}^{2} \Big(\nu_i^I(Q) \otimes \nu_i^{II}(P) - \nu_i^I(P) \otimes \nu_i^{II}(Q)\Big), \end{aligned} \tag{3.11}$$

where

$$d_Q\Sigma\big(P, Q\big) := dx_P \otimes dx_Q \frac{\partial}{\partial x_Q} \frac{y_{4,P}y_{5,P} + y_{4,P}y_{5,Q} + y_{4,Q}y_{5,P}}{(x_P - x_Q)3y_{4,P}y_{5,P}}. \tag{3.12}$$

The set of differentials $\{\nu_1^{II}, \nu_2^{II}\}$ *is determined modulo the* $\mathbb{C}$*-linear space spanned by* $\langle \nu_j^I \rangle_{j=1,2}$ *and the components of its representative one are*

$$\{\nu_1^{II}, \nu_2^{II}\} = \left\{ \frac{-\left(2x + \lambda_1^{(2)}\right) dx}{3y_4} . \frac{-xdx}{3y_5} \right\}.$$

We will fix this ν_i^{II} *hereafter.*

Proof.

$$\frac{\partial}{\partial x_Q} \frac{y_{4,P}y_{5,P} + y_{4,P}y_{5,Q} + y_{4,Q}y_{5,P}}{(x_P - x_Q)3y_{4,P}y_{5,P}} dx_P$$
$$= \frac{1}{(x_P - x_Q)9y_{4,P}y_{5,P}y_{4,Q}y_{5,Q}} \Big[\frac{3(y_{4,P}y_{5,P} + y_{4,P}y_{5,Q} + y_{4,Q}y_{5,P})y_{4,Q}y_{5,Q}}{(x_P - x_Q)}$$
$$+ \Big(y_{4,P} \frac{y_{4,Q}}{y_{5,Q}} (2k_{2,Q}k'_{2,Q}k_{1,Q} + k_{2,Q}^2 k'_{1,Q}) + y_{5,P} \frac{y_{5,Q}}{y_{4,Q}} (2k_{2,Q}k_{1,Q}k'_{1,Q} + k'_{2,Q}k_{1,Q}^2)) \Big) \Big].$$

Here $k_{a,P} = k_a(x_P)$ and $k'_{a,P} = dk_a(x_P)/dx_P$.

$$\frac{\partial}{\partial x_Q} \frac{y_{4,P}y_{5,P} + y_{4,P}y_{5,Q} + y_{4,Q}y_{5,P}}{(x_P - x_Q)3y_{4,P}y_{5,P}} - \frac{\partial}{\partial x_P} \frac{y_{4,Q}y_{5,Q} + y_{4,Q}y_{5,P} + y_{4,P}y_{5,Q}}{(x_Q - x_P)3y_{4,Q}y_{5,Q}}$$

is equal to

$$= \frac{1}{(x_P - x_Q)9y_{4,P}y_{5,P}y_{4,Q}y_{5,Q}} \Big[\frac{3(y_{4,P}y_{5,Q} + y_{4,Q}y_{5,P})(y_{4,Q}y_{5,Q} - y_{4,P}y_{5,P})}{(x_P - x_Q)}$$
$$- \Big(y_{4,P}y_{5,Q}(2k'_{2,Q}k_{1,Q} + k_{2,Q}k'_{1,Q} - 2k_{2,P}k'_{1,P} + k'_{2,P}k_{1,P})$$
$$+ y_{4,Q}y_{5,P}(2k'_{2,P}k_{1,P} + k_{2,P}k'_{1,P} - 2k_{2,Q}k'_{1,Q} + k'_{2Q}k_{1,Q}) \Big) \Big]$$
$$= \frac{1}{(x_P - x_Q)9y_{4,P}y_{5,P}y_{4,Q}y_{5,Q}} \left(B_2(P,Q) - B_2(Q,P) \right).$$

where

$$B_2(P,Q) = y_{4,P}y_{5,Q}\left(2x_Q + \lambda_1^{(2)} - x_P\right).$$

Then we have the result. □

Corollary 3.7. *(1) The one form,*

$$\Pi_{P_1}^{P_2}(P) := \Sigma(P, P_1) - \Sigma(P, P_2),$$

is a differential of the third kind, whose only (first-order) poles are $P = P_1$ *and* $P = P_2$*, and residues* $+1$ *and* -1 *respectively.*

(2) The fundamental differential of the second kind $\Omega(P_1, P_2)$ is given by

$$\begin{aligned}\Omega(P_1, P_2) &= d_{P_2}\Sigma(P_1, P_2) + \sum_{i=1}^{2} \nu^I_i(P_1) \otimes \nu^{II}_i(P_2) \\ &= \frac{F(P_1, P_2) dx_1 \otimes dx_2}{(x_1 - x_2)^2 9 y_{4,P_1} y_{5,P_1} y_{4,P_2} y_{5,P_2}},\end{aligned} \tag{3.13}$$

where F is an element of $R \otimes R$.

Proof. Direct computations show it. □

Lemma 3.8. *We have*

$$\lim_{P_1 \to \infty} \frac{F(P_1, P_2)}{\phi_{{H^1}_1}(P_1)(x_1 - x_2)^2} = \phi_{{H^1}_2}(P_2) = x_{P_2} y_{4,P_2}. \tag{3.14}$$

Proof. B_2 in the proof of Proposition 3.6 shows the result. □

For later convenience we introduce the quantity:

$$\begin{aligned}\Omega^{P_1,P_2}_{Q_1,Q_2} &:= \int_{P_2}^{P_1} \int_{Q_2}^{Q_1} \Omega(P, Q) \\ &= \int_{P_2}^{P_1} (\Sigma(P, Q_1) - \Sigma(P, Q_2)) + \sum_{i=1}^{4} \int_{P_2}^{P_1} \nu^I_i(P) \int_{Q_2}^{Q_1} \nu^{II}_i(P).\end{aligned} \tag{3.15}$$

4. THE SIGMA FUNCTION OVER $(3, 4, 5)$ CURVE

4.1. **Legendre relation.** Corresponding to (3.6), we define the complete integral of the second kind,

$$[\eta' \ \eta''] := \frac{1}{2} \left[\int_{\alpha_i} \nu^{II}_j \quad \int_{\beta_i} \nu^{II}_j \right]_{i,j=1,2}. \tag{4.1}$$

Let τ_{Q_1,Q_2} be the normalized differential of the third kind such that τ_{Q_1,Q_2} has residues $+1$, -1 at Q_1 and Q_2, is regular everywhere else, and is normalized, $\int_{\alpha_i} \tau_{P,Q} = 0$ for $i = 1, 2$ [FK, III.3.5]. The following Lemma corresponding to Corollary 2.6 (ii) in [F] holds.

Lemma 4.1. *By letting $\gamma = {\omega'}^{-1}\eta'$, we have*

$$\Omega^{P_1,P_2}_{Q_1,Q_2} = \int_{P_2}^{P_1} \tau_{Q_1,Q_2} + \sum_{i,j=1}^{2} \gamma_{ij} \int_{P_2}^{P_1} \nu^I_i \int_{Q_2}^{Q_1} \nu^I_j.$$

Proof. We note that $\partial_{P_1}\Omega^{P_1,P_2}_{Q_1,Q_2} - \tau_{Q_1,Q_2}$ is expressed by a linear combination of the holomorphic one forms. Noting $\int_{P_1}^{P_2}\tau_{Q_1,Q_2} = \int_{Q_1}^{Q_2}\tau_{P_1,P_2}$ [FK, III.3.5], γ_{ij} is symmetric. Due to (3.15),

$$(\Sigma(P_1,Q_1)-\Sigma(P_1,Q_2)) + \sum_{i=1}^{g}\nu_i^I(P_1)\int_{Q_2}^{Q_1}\nu_i^{II}(P) = \tau_{Q_1,Q_2}(P_1) + \sum_{i,j=1}^{2}\gamma_{ij}\nu_i^I(P_1)\int_{Q_2}^{Q_1}\nu_j^I.$$

By choosing an appropriate path Γ from Q_1 to Q_2, homotopic to α_k, we have $(\eta'\nu^I)^t = \gamma\cdot(\omega'\nu^I)^t$. ☐

The following Proposition provides a symplectic structure in the Jacobian $\mathcal{J}_2$, known as *generalized Legendre relation* [B1, BLE1, BLE2].

Proposition 4.2. (generalized Legendre relation) *The matrix,*

$$M := \begin{bmatrix} 2\omega' & 2\omega'' \\ 2\eta' & 2\eta'' \end{bmatrix}, \tag{4.2}$$

satisfies

$$M\begin{bmatrix} & -1 \\ 1 & \end{bmatrix}{}^tM = 2\pi\sqrt{-1}\begin{bmatrix} & -1 \\ 1 & \end{bmatrix}. \tag{4.3}$$

Proof. For example, for

$$\int_{P_2}^{P_1}(\Sigma(P,Q_1)-\Sigma(P,Q_2)) + \sum_{i=1}^{g}\int_{P_2}^{P_1}\nu_i^I(P)\int_{Q_2}^{Q_1}\nu_i^{II}(P)$$
$$= \int_{Q_2}^{Q_1}\tau_{P_1,P_2} + \sum_{i,j=1}^{2}\gamma_{ij}\int_{P_2}^{P_1}\nu_i^I\int_{Q_2}^{Q_1}\nu_j^I,$$

the contour of Q_1 to Q_2 along α_k gives

$$\sum_{i=1}^{2}\int_{P_2}^{P_1}\nu_i^I(P)\eta'_{ki} = \sum_{i,j=1}^{2}\gamma_{ij}\int_{P_2}^{P_1}\nu_i^I\omega'_{kj},$$

and further contour integration provides ${}^t({}^t\omega'\eta') = ({}^t\omega'\eta')$. By considering the contour integral, the contour of Q_1 to Q_2 along β_k gives

$$\sum_{i=1}^{2}\int_{P_2}^{P_1}\nu_i^I(P)\eta''_{ki} = 2\pi\sqrt{-1}\int_{P_2}^{P_1}(\omega'^{-1}\nu^I)_k + \sum_{i,j=1}^{2}\gamma_{ij}\int_{P_2}^{P_1}\nu_i^I\omega'_{kj},$$

which turns out to be

$${}^t({}^t\omega''\eta'') = 2\pi\sqrt{-1}\mathbb{T} + ({}^t\omega''\gamma\omega''),$$
$$({}^t\omega'\eta'') = 2\pi\sqrt{-1}1_2 + ({}^t\omega'\gamma\omega'').$$

Here we use $\int_{\beta_k}\tau_{Q_1,Q_2} = 2\pi\sqrt{-1}\int_{Q_2}^{Q_1}(\omega'^{-1}\nu^I)_k$ [FK, III.3.5]. ☐

4.2. σ **function.** Due to the Riemann relations [F], $\mathrm{Im}\,(\omega'^{-1}\omega'')$ is positive definite. Theorem 1.1 in [F] gives

$$\delta := \begin{bmatrix} \delta' \\ \delta'' \end{bmatrix} \in \left(\frac{\mathbb{Z}}{2}\right)^4 \tag{4.4}$$

be the theta characteristic which is equal to the Riemann constant ξ_R with respect to the base point ∞ and the period matrix $[\,2\omega'\ 2\omega'']$.

We define an entire function of (a column-vector) $u = {}^t(u_1, u_2) \in \mathbb{C}^2$,

$$\begin{aligned} \sigma(u) &= \sigma(u; M) = \sigma(u_1, u_2; M) \\ &= c\exp(-\tfrac{1}{2}\ {}^t u\eta'\omega'^{-1}u)\vartheta[\delta]\,(\frac{1}{2}\omega'^{-1}u;\ \omega'^{-1}\omega'') \\ &= c\exp(-\tfrac{1}{2}\ {}^t u\eta'\omega'^{-1}\ u) \\ &\qquad\times \sum_{n\in\mathbb{Z}^2} \exp\left[\pi\sqrt{-1}\{\ {}^t(n+\delta')\omega'^{-1}\omega''(n+\delta') +\ {}^t(n+\delta')(\omega'^{-1}u+\delta'')\}\right], \end{aligned} \tag{4.5}$$

where c is a certain constant of a rational of b's.

For a given $u \in \mathbb{C}^2$, we introduce u' and u'' in $\mathbb{R}^2$ so that

$$u = 2\omega' u' + 2\omega'' u''.$$

Proposition 4.3. *For $u,\ v \in \mathbb{C}^2$, and $\ell\ (= 2\omega'\ell' + 2\omega''\ell'') \in \Pi_2$, we define*

$$\begin{aligned} L(u, v) &:= 2\ {}^t u(\eta' v' + \eta'' v''), \\ \chi(\ell) &:= \exp[\pi\sqrt{-1}\big(2({}^t\ell'\delta'' - {}^t\ell''\delta') + {}^t\ell'\ell''\big)]\ (\in \{1, -1\}). \end{aligned}$$

The following holds

$$\sigma(u+\ell) = \sigma(u)\exp(L(u+\frac{1}{2}\ell, \ell))\chi(\ell). \tag{4.6}$$

Proof. It is essentially the same as one for the normalized theta function [L, Chap. VI]. □

The vanishing locus of σ is simply given by

$$\Theta^1 := (\mathcal{W}^1 \cup [-1]\mathcal{W}^1) = \mathcal{W}^1. \tag{4.7}$$

4.3. **The Riemann fundamental relation.** We review a relation which we call the Riemann fundamental relation [B1, §195]:

Proposition 4.4. *For $(P, Q, P_i, P_i') \in X^2 \times (S^2(X) \setminus S_1^2(X)) \times (S^2(X) \setminus S_1^2(X))$,*

$$u := \sum_{i=1}^{2} w(P_i), \quad v := \sum_{i=1}^{2} w(P_i'),$$

$$\begin{aligned}\exp\left(\sum_{i,j=1}^{2}\Omega_{P_i,P'_j}^{P,Q}\right) &= \frac{\sigma(w(P)-u)\sigma(w(Q)-v)}{\sigma(w(Q)-u)\sigma(w(P)-v)}\\ &= \frac{\sigma(w(P)-w(P_1,P_2))\sigma(w(Q)-w(P'_1,P'_2))}{\sigma((w(Q)-w(P_1,P_2))\sigma(w(P)-w(P'_1,P'_2))}.\end{aligned}$$

Proof. The right-hand side is expressed by

$$\exp(\text{bilinear term in } w\text{'s})\frac{\theta(\omega'^{-1}(w(P)-u)+\xi_R)\theta(\omega'^{-1}(w(Q)-v)+\xi_R)}{\theta(\omega'^{-1}(w(Q)-u)+\xi_R)\theta(\omega'^{-1}(w(P)-v))+\xi_R)},$$

where ξ_R is the Riemann constant. By Riemann's theorem for theta functions [F, p.23], the above becomes

$$\exp(\text{bilinear term in } w\text{'s})\exp\left(\sum_{j=1}^{2}\int_Q^P \tau_{P_j,P'_j}\right).$$

The exponential part of the bilinear term is given by,

$$(u-v)^t\gamma(w(P)-w(Q)) = \sum_{i,j,k}\gamma_{ij}\int_{P'_k}^{P_k}\nu_i^I\int_Q^P\nu_j^I,$$

which equals to (Lemma (4.1))

$$\sum_{i=1}^{2}\Omega_{P_i,P'_i}^{P,Q} - \sum_{i=1}^{2}\int_Q^P \tau_{P_i,P'_i}.$$

The above integrals depend upon the paths we choose, but due to (4.6) such dependence cancels. We have the equality. □

Proposition 4.5. *For* $(P,P_1,P_2)\in X\times S^2(X)\setminus S^2_1(X)$ *and* $u := w(P_1,P_2)$*, the equality*

$$\sum_{i,j=1}^{2}\wp_{i,j}\left(w(P)-u\right)\phi_{H^1{}_{i-1}}(P)\phi_{H^1{}_{j-1}}(P_a) = \frac{F(P,P_a)}{(x-x_a)^2},$$

holds for every $a=1,2$*, where we set*

$$\wp_{ij}(u) := -\frac{\sigma_i(u)\sigma_j(u)-\sigma(u)\sigma_{ij}(u)}{\sigma(u)^2} \equiv -\frac{\partial^2}{\partial u_i\partial u_j}\log\sigma(u).$$

Proof. For the case X, using the relation

$$\sum_{i,j=1}^{2}\phi_{H^1{}_{i-1}}(P_1)\phi_{H^1{}_{j-1}}(P_2)\frac{\partial^2}{\partial w_i(P_1)\partial w_j(P_2)} = 9y_{4,P_1}y_{5,P_1}y_{4,P_2}y_{5,P_2}\frac{\partial^2}{\partial x_1\partial x_2},$$

taking logarithm of both sides of the relation in Proposition 4.4 and differentiating them along $P_1=P$ and $P_2=P_a$, we obtain the claim. □

4.4. **The μ_n functions over** $(3,4,5)$**.** As mentioned in [MP08], we introduce meromorphic functions on the curve X.

For the definition of the function μ's, we introduce the Frobenius-Stickelberger (FS) matrix and its determinant following [MP08]. Let n be a positive integer and $P_1, \dots, P_n$ be in $X \backslash \infty$. We define the *Frobenius-Stickelberger* (FS) *matrix* by:

$$\Psi_n(P_1, P_2, \dots, P_n) := \begin{pmatrix} \phi_{H^1 0}(P_1) & \phi_{H^1 1}(P_1) & \phi_{H^1 2}(P_1) & \cdots & \phi_{H^1 n-1}(P_1) \\ \phi_{H^1 0}(P_2) & \phi_{H^1 1}(P_2) & \phi_{H^1 2}(P_2) & \cdots & \phi_{H^1 n-1}(P_2) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \phi_{H^1 0}(P_n) & \phi_{H^1 1}(P_n) & \phi_{H^1 2}(P_n) & \cdots & \phi_{H^1 n-1}(P_n) \end{pmatrix}.$$

and $\psi_n(P_1, P_2, \dots, P_n) := \det \Psi_n(P_1, P_2, \dots, P_n)$. We call this matrix *Frobenius-Stickelberger (FS) matrix* and its determinant *Frobenius-Stickelberger (FS) determinant.* These become singular for some tuples in $(X \backslash \infty)^n$.

Definition 4.6. *For* $P, P_1, \dots, P_n \in (X \backslash \infty) \times SS^n(X \backslash \infty)$*, we define* $\mu_n(P)$ *by*

$$\mu_n(P) := \mu_n(P; P_1, \dots, P_n) := \lim_{P'_i \to P_i} \frac{1}{\psi_n(P'_1, \dots, P'_n)} \psi_{n+1}(P'_1, \dots, P'_n, P),$$

where the P'_i *are generic, the limit is taken (irrespective of the order) for each* i*; and* $\mu_{n,k}(P_1, \dots, P_n)$ *by*

$$\mu_n(P) = \phi_{H^1 n}(P) + \sum_{k=0}^{n-1} (-1)^{n-k} \mu_{n,k}(P_1, \dots, P_n) \phi_{H^1 k}(P).$$

For example,

$$\begin{aligned} \mu_2(P; P_1, P_2) &= x y_4 - \frac{y_{4,1} x_2 y_{4,2} - y_{4,2} x_1 y_{4,1}}{y_{4,1} y_{4,2} - y_{4,2} y_{4,1}} y_5 + \frac{y_{5,1} x_2 y_{4,2} - y_{5,2} x_1 y_{4,1}}{y_{4,1} y_{4,2} - y_{4,2} y_{4,1}} y_4, \\ \mu_1(P; P_1) &= y_5 + \frac{y_{5,1}}{y_{4,1}} y_4. \end{aligned}$$

We note that $\mu_{H^1,n}$ for X is characterized by the condition on a polynomial $\alpha_{H^1 n} = \sum_{i=0}^{n} a_i \phi_{H^1 i}(P)$, $a_i \in \mathbb{C}$ and $a_n = 1$, which has a zero at each point P_i and has the smallest possible order such that it multiplied by $dx/3y_4y_5$ belongs to $H^1(X \setminus \infty, \mathcal{O}_X)$.

4.5. **Jacobi inversion formulae over Θ^k.**

Theorem 4.7. (Jacobi inversion formula) *For* $(P, P_1, P_2) \in X \times (S^2(X) \setminus S^2_1(X))$*, we have*

(1) $\mu_{H^1,2}(P; P_1, P_2) = x y_4 - \wp_{22}(w(P_1, P_2)) y_4 + \wp_{21}(w(P_1, P_2)) y_5$.

(2) $\wp_{22}(w(P_1, P_2)) = \dfrac{y_{4,1} x_2 y_{4,2} - y_{4,2} x_1 y_{4,1}}{y_{4,1} y_{4,2} - y_{4,2} y_{4,1}}$, $\wp_{21}(w(P_1, P_2)) = \dfrac{y_{5,1} x_2 y_{4,2} - y_{5,2} x_1 y_{4,1}}{y_{4,1} y_{4,2} - y_{4,2} y_{4,1}}$.

Proof. the same as [MP08]. □

Theorem 4.8. *For*

$$\Theta^1 := \mathcal{W}^1 = [-1]\mathcal{W}^1, \tag{4.8}$$

and $P_1 \in X \setminus S_1^1(X)$ *and* $u = \pm w(P_1) \in \kappa^{-1}(\Theta^1)$,

$$\frac{\sigma_1(u)}{\sigma_2(u)} = \frac{y_5}{y_4}$$

Proof. As in [MP08], by considering $\lim_{P_2 \to \infty} \dfrac{\wp_{21}(P_1, P_2)}{\wp_{22}(P_1, P_2)}$, we have the result. □

Following the conjecture by Buchstaber, Leykin and Enolskii [BEL2], in [Na] Nakayashiki showed that the leading of the σ function for (r, s) curve is expressed by Schur function. Noting (3.5) and (3.7), the above Jacobi inversion formulae gives an extension that

$$\sigma(u) = \frac{1}{2}u_2^2 - u_1 + \sum_{|\alpha|>2} a_\alpha u^\alpha,$$

where $a_\alpha \in \mathbb{Q}[b_1, \cdots, b_5]$, $\alpha = (\alpha_1, \alpha_2)$, $|\alpha| = \alpha_1 + \alpha_2$ and $u^\alpha = u_1^{\alpha_1} u_2^{\alpha_2}$. Since for a Young diagram Λ, S_Λ and s_Λ are the Schur functions defined by

$$S_\Lambda(T_1, T_2) = s_\Lambda(t_1, t_2) = t_1 t_2 = \frac{1}{2}T_1^2 - T_2,$$

where $T_1 := t_1 + t_2$ and $T_2 := \frac{1}{2}(t_1^2 + t_2^2)$, we have

$$\sigma(u) = S_\Lambda(u_1, u_2) + \sum_{|\alpha|>2} a_\alpha u^\alpha.$$

5. DISCUSSION

We showed that the EEL construction works well even for a space curve, and the sigma function is naturally defined. More precisely we applied the EEL construction to the curve X of $(3, 4, 5)$ type and obtained a fundamental differential of the second kind, natural differentials of the second kind, and the third kind. Using them we define the sigma function even for the space curve X explicitly. Further the sigma function gives the Jacobi inversion formulae for the Jacobian $\mathcal{J}$ and even for strata in $\mathcal{J}$ as a natural extension of [MP08].

Since this construction is very natural, this study sheds a new light on the way to construction of the sigma functions for space curves. We conjectured that the EEL construction could be applied to every space curve if it is Weierstrass. In fact, it is applicable for the curves, $C(H_4)$ and $C(H_{12})$ generated by $\langle 3, 7, 8\rangle$ and $\langle 6, 13, 14, 15, 16\rangle$ [KMP].

As mentioned in Introduction, degenerate curves naturally appear when we deal with (r, s) curves in a physical problem e.g., [Ma08], the degenerate curves might be space curves by normalisation. This construction is applied to a more interesting physical phenomenon.

As an interesting example of a space curve, we will give a comment on a problem as follows, for which I started to study sigma functions for affine space curves.

McKay considers a relation between dispersionless KP hierarchy and the replicable functions in order to obtain a further profound interpretation of the moonshine phenomena of Monster group [FMN, Mc, MS]. He conjectured that it might be related to the quantised elastica [Ma08]. By studying a relation between a replicable function and an algebraic curve associated with elastica, we found that a semigroup H_{12} generated by $M_{12} := \langle 6, 13, 14, 15, 16 \rangle$ has gap sequence,

$$L(H_{12}) = \{1, 2, 3, 4, 5, 7, 8, 9, 10, 11, 17, 23\},$$

which is identical to the Norton number,

$$N_{12} := \{1, 2, 3, 4, 5, 7, 8, 9, 11, 17, 19, 23\}$$

by exchanging 10 and 19. The Norton number plays the essential role in the moonshine phenomena for the Monster group [Cu, N]. Norton showed that the replicable function is given as an element of $\mathbb{Q}[a_1, a_2, a_3, a_4, a_5, a_7, a_8, a_9, a_{11}, a_{17}, a_{19}, a_{23}][[t]]$ [Cu, FMN, N]. The replicable function is a generalization of the elliptic J-function, which causes the moonshine phenomena of the Monster group.

After then, Komeda proved that H_{12} is the Weierstrass semigroup by constructing a non-singular algebraic curve $C(H_{12})$ using a fibre structure over a curve $C(H_4)$, as mentioned in Appendix. He also gave the fundamental relations in Propositions A.1 and A.2 which are reported in [KMP].

In [KMP], using the results, we apply the EEL construction to the curves $C(H_4)$ and $C(H_{12})$, and obtain a fundamental differential of the second kind and natural differentials of the second kind, and the third kind. Using them, we also define the sigma functions even for the space curves $C(H_6)$ and $C(H_{12})$ explicitly which give the Jacobi inversion formulae. The numbers 10 and 19 are related to the weight of the denominator of the holomorphic one forms in $C(H_{12})$. The sigma function is related to the Schur function $S_{\Lambda_{12}}(T_1, T_2, T_3, T_4, T_5, T_7, T_8, T_9, T_{10}, T_{11}, T_{17}, T_{23})$ for the Young diagram;



.

Since the component u_i of the element u in the Jacobi variety $\mathcal{J}_{12}$ for $C(H_{12})$ has a natural weight $w(u_i)$ as in (3.7), u can be expressed by $u = (v_{23}, v_{19}, v_{17}, v_{11}, v_9, v_8, v_7, v_5, v_4, v_3, v_2, v_1)$ by denoting $v_{w(u_i)} = u_i$ and then v_j can be regarded as Sato coordinate in the KP hierarchy. Then σ is a function of $(v_1, v_2, v_3, v_4, v_5, v_7, v_8, v_9, v_{11}, v_{17}, v_{19}, v_{23})$. If we consider a deformation of $C(H_{12})$ parameterized by $q \in \mathbb{C}$, a truncated part of the expansion of the sigma function belongs to $\mathbb{Q}[v_1, v_2, v_3, v_4, v_5, v_7, v_8, v_9, v_{11}, v_{17}, v_{19}, v_{23}][q]$, whose properties are similar to the replicable functions [KMP].

Since the Jacobi variety $\mathcal{J}_{12}$ for $C(H_{12})$ is given as 12-dimensional complex torus whose real dimension is 24. Hence it might remind us of Witten conjecture associated with Monster group problem [HBJ]; Witten conjectured that a 24 dimensional manifold exists such that the Monster group acts on it.

Further $C(H_{12})$ naturally contains a sextic curve, which is related to $\mathbb{E}_8$ [Co]. In a subvariety $\mathcal{W}^k$ of $\mathcal{J}_{12}$ $(k < 12)$, v_j $(j = w(u_i); i = k+1, k+2, \dots, 12)$ is a function of $v_{j'}$, $(j' = w(u_i); i = 1, 2, \dots, k)$ is a function of $v_{j'}$, σ function has a reduction such that $\sigma_{\natural k}$ is a function related to $\mathcal{W}^k$ [MP11]. It also reminds us of the fact that $a_5, \dots, a_{23}$ are functions of a_1, a_2, a_3 and a_4 in the elliptic J-function as a replicable function.

A. APPENDIX: WEIERSTRASS PROPERTIES OF $(6, 13, 14, 15, 16)$ BY JIRYO KOMEDA
[1]

Proposition A.1. *The numerical semigroup* $\langle 6, 13, 14, 15, 16 \rangle$ *is Weierstrass.*

This proof is given in [KMP] but we show its sketch.

Proof. Let (C, P) be a pointed curve with $H(P) = \langle 3, 7, 8 \rangle$. Then

$$2 = h^0(4P) = 4 + 1 - 4 + h^0(K - 4P) = 1 + h^0(K - 4P),$$

which implies that $K - 4P \sim P_1 + P_2$ for some points P_1 and $P_2 \in C$. Here K is a canonical divisor on C. Moreover,

$$2 = h^0(5P) = 5 + 1 - 4 + h^0(K - 5P) = 2 + h^0(K - 5P),$$

which implies that $h^0(K - 5P) = 0$. Hence, we get $P_i \neq P$ for $i = 1, 2$. Thus, $K \sim 4P + P_1 + P_2$ with $P_i \neq P$ for $i = 1, 2$. We set $D = 7P - P_1 - P_2$. Then $\deg(2D - P) = 9 = 2 \times 4 + 1$, which implies that the complete linear system $|2D - P|$ is very ample, hence base-point free. Therefore, $2D \sim P + Q_1 + \ldots + Q_9$ $(=$ a reduced divisor$)$. Let $\mathcal{L}$ be the invertible sheaf $\mathcal{O}_C(-D)$ on C and ϕ an isomorphism $\mathcal{L}^{\otimes 2} \approx \mathcal{O}_C(-P - Q_1 - \cdots - Q_9) \subset \mathcal{O}_C$. Then the vector bundle $\mathcal{O}_C \oplus \mathcal{L}$ has an $\mathcal{O}_C$-algebra structure through ϕ. The canonical morphism $\pi : \tilde{C} = \mathrm{Spec}\ (\mathcal{O}_C \oplus \mathcal{L}) \to C$, is a double covering. Its branch locus of π is

---

[1] Department of Mathematics
Center for Basic Education and Integrated Learning,
Kanagawa Institute of Technology,
Atsugi, 243-0292, JAPAN.
e-mail: komeda@gen.kanagawa-it.ac.jp

$\{P, Q_1, ..., Q_9\}$. Let $\tilde{P}$ be the ramification point of π over P. Then it can be showed that $H(\tilde{P}) = \langle 6, 13, 14, 15, 16\rangle$ using the formula, $h^0(2n\tilde{P}) = h^0(nP) + h^0(nP - D)$ for any non-negative integer n.

By considering $h^0(2n\tilde{P})$ for $n = 3, 4, 5, 6, 7, 8, 9$, we show $H(\tilde{P}) = \langle 6, 13, 14, 15, 16\rangle$. $\square$

Further we have the following proposition whose proof is in [KMP]:

Proposition A.2. *Let* B_{12} *a monomial ring which is given by* $k[t^a]_{a\in M_{12}}$ *for the numerical semigroup* H_{12}. *For a* k*-algebra homomorphism,*

$$\varphi_{12} : k[Z] := k[Z_6, Z_{13}, Z_{14}, Z_{15}, Z_{16}] \to k[t^a]_{a\in M_{12}}$$

where Z_a *is the weight of* $a = 6, 13, 14, 15, 16$, *the kernel of* φ_{12} *is generated by the following relations* $f^{(Z)}_{12,b}$ $(b = 1, \cdots, 9)$,

$$\begin{array}{lll} f^{(Z)}_{12,1} = Z_{13}^2 - Z_6^2 Z_{14}, & f^{(Z)}_{12,2} = Z_{13}Z_{14} - Z_6^2 Z_{15}, & f^{(Z)}_{12,3} = Z_{14}^2 - Z_{13}Z_{15}, \\ f^{(Z)}_{12,4} = Z_{14}^2 - Z_6^2 Z_{16}, & f^{(Z)}_{12,5} = Z_{13}Z_{16} - Z_{14}Z_{15}, & f^{(Z)}_{12,6} = Z_{15}^2 - Z_6^5, \\ f^{(Z)}_{12,7} = Z_{14}Z_{16} - Z_6^5, & f^{(Z)}_{12,8} = Z_{15}Z_{16} - Z_6^3 Z_{13}, & f^{(Z)}_{12,9} = Z_{16}^2 - Z_6^3 Z_{14}. \end{array} \tag{A.1}$$

References

[ACGH] E. Arbarello, M. Cornalba, P.A. Griffiths and J. Harris, Geometry of Algebraic Curves. Vol. I, Springer-Verlag, New York, 1985.

[B1] H.F. Baker, Abelian functions. Abel's theorem and the allied theory of theta functions, Cambridge University Press, Cambridge, 1995, Reprint of the 1897 original.

[B2] H.F. Baker, *On the hyperelliptic sigma functions*, Amer. J. Math., **20** (1898) 301-384.

[B3] H.F. Baker, *On a system of differential equations leading to periodic functions*, Acta Math., **27** (1903) 135-156.

[BG] S. Baldwin and J. Gibbons, *Genus 4 trigonal reduction of the Benney equations*, J. Phys. A: Math. Gen., **39** (2006) 3607-3639.

[BDE] H. W. Braden, D. Antonella and V.Z. Enolski, *On charge-3 cyclic monopoles*, Nonlinearity, **24** (2011) 643-675.

[BEL1] V.M. Buchstaber, V.Z. Enolskiĭ and D.V. Leĭkin, *Hyperelliptic Kleinian functions and applications*, in Solitons, geometry, and topology: on the crossroad, 1–33, Amer. Math. Soc. Transl. Ser. 2, 179, Amer. Math. Soc., Providence, RI, 1997.

[BEL2] V.M. Bukhshtaber, V.Z. Ènol'skiĭ and D.V. Leĭkin, *Rational analogues of Abelian functions*, Funct. Anal. Appl., **33** (1999) 83-94.

[BLE1] V.M. Bukhshtaber, D.V. Leĭkin, and V.Z. Ènol'skiĭ, *σ-functions of (n, s)-curves*, Russian Math. Surveys, **54** (1999) 628-629.

[BLE2] V.M. Bukhstaber, D.V. Leĭkin, and V.Z. Ènol'skiĭ, *Uniformization of Jacobi manifolds of trigonal curves and nonlinear differential equations*, Funct. Anal. Appl., **34** (2000) 159-171.

[Co] A. B. Coble, Algebraic geometry and theta functions, Amer. Math. Soc., New York, 1961, .

[Cu] C. J. Cummins, *Some comments on replicable functions*, Modern trends in Lie algebra representation theory, (Queen's Univ., Kingston, ON, 1994) 48-55, Queen's paper in Pure and Appl. Math **94** (1994) .

[EEL] J.C. Eilbeck, V.Z. Enolskii and D.V. Leykin, *On the Kleinian construction of abelian functions of canonical algebraic curves*, SIDE III—symmetries and integrability of difference equations (Sabaudia, 1998), 121–138, CRM Proc. Lecture Notes, 25, Amer. Math. Soc., Providence, RI, 2000.

[EEMOP1] J.C. Eilbeck, V.Z. Enolskii, S. Matsutani, Y. Ônishi, E. Previato, *Addition formulae over the Jacobian preimage of hyperelliptic Wirtinger varieties*, J. reine angew. Math., **619** (2008) 37–48.

[EEMOP2] J.C. Eilbeck, V.Z. Enolskii, S. Matsutani, Y. Ônishi, E. Previato, *Ableian Functions for Trigonal Curves of Genus Three*, Int. Math. Res. Not., **2007** (2007) 140,38.

[FK] H. M. Farkas and I. Kra, Riemann Surfaces, second edition, Graduate Texts in Mathematics, Vol. 71. Springer-Verlag, Berlin-New York, 1991.

[F] J.D. Fay, Theta functions on Riemann Surfaces, Lecture Notes in Mathematics, Vol. 352. Springer-Verlag, Berlin-New York, 1973.

[FMN] D. J. Ford, J. McKay and S. P. Norton, *More on replicable functions*, Comm. in Alg., **22** (1994) 5175-5193.

[HBJ] F. Hirzebruch, T. Berger and R. Jung, Manifolds and Modular Forms, Vieweg, Bonn, 1992.

[Gr] D. Grand, *A generalization of a formula of Eisenstein*, Proc. Lond. Math. Soc., **62** (1991) 121–132.

[K] F. Klein, *Über hyperelliptische Sigmafunctionen*, Math. Ann., **32** (1888) 351-380.

[K83] J. Komeda, *On Weierstrass points whose first non-gaps are four*, J. Reine. Angew. Math., **341** (1983), 68-86.

[K99] J. Komeda, *Existence of the primitive Weierstrass gap sequences on curves of genus 9*, Bol. Soc. Bras. Mat., **30** (1999), 125-137.

[K04] J. Komeda, *On 6-semigroups generated by 4-elements from which affine toric varieties can be constructed*, Research Reports of Kanagawa Institute of Technology **B-28** (2004), 79-85.

[KO04] J. Komeda and A. Ohbuchi *Weierstrass points with first non-gap four on a double covering of a hyperelliptic curve*, Serdica Math J., **30** (2004), 43-54; Corrigendum **32** (2006), 375-378.

[KO08] J. Komeda and A. Ohbuchi *Weierstrass points with first non-gap four on a double covering of a hyperelliptic curve II*, Serdica Math J., **34** (2008), 771-782.

[KO08a] J. Komeda and A. Ohbuchi *Existence of the non-primitive Weierstrass gap sequences on curves of genus 8*, Bull Braz Math. Soc. News Series, **39** (2008), 109-121.

[KMP] J. Komeda, S. Matsutani, E. Previato *Sigma functions over space curves* $(3,7,8)$ *and* $(6,13,14,15,16)$, in preparation, .

[L] S. Lang, Introduction to algebraic and abelian functions, Second edition, Graduate Texts in Mathematics, 89. Springer-Verlag, New York-Berlin, 1982.

[Mat] H. Matsumura, Commutative ring theory, CSAM 8, , translated by M. Reid, Cambridge, Cambridge, 1986.

[Ma01] S. Matsutani, *Hyperelliptic solutions of KdV and KP equations: re-evaluation of Baker's study on hyperelliptic sigma functions*, J. Phys. A, **34** (2001) 4721–4732.

[Ma08] S. Matsutani, *Relations in a quantized elastica*, J. Phys. A, **41** (2008) 075201.

[MP08] S. Matsutani and E. Previato, Jacobi inversion on strata of the Jacobian of the C_{rs} curve $y^r = f(x)$, J. Math. Soc. Japan **60** (2008) 1009-1044.

[MP09] S. Matsutani and E. Previato, A class of solutions of the dispersionless KP equation, Phys. Lett., (2009) .

[MP11] S. Matsutani and E. Previato, Jacobi inversion on strata of the Jacobian of the C_{rs} curve $y^r = f(x)II$, in preparation.

[Mc] J. McKay, *Essentials of Monstrous Moonshine*, Adv. Std. Pure Math., **32** (2001) 347-353.

[MS] J. McKay, and A. Sebbar, Fuchsian groups, automorphic forms and Schwarzians, Math. Ann. 318 (2000), no. 2, 255-275.

[Mu] D. Mumford, Tata Lectures on Theta II Jacobian theta functions and differential equations, Progress in Mathematics, 43. Birkhäuser Boston, Inc., Boston, MA, 1984.

[Na] A. Nakayashiki, *On algebraic expansions of sigma functions for* (n, s) *curves*, arXiv:0803.2083.

[N] S. P. Norton More on Moonshine, in Computational Group Theory eded M D. Atkinson, 1984 Academic Press, p.185-194.

[O] Y. Ônishi, *Determinant expressions for hyperelliptic functions*, Proc. Edinb. Math. Soc., **48** (2005) 1-38.

[P] H. C. Pinkham, *Deformation of algebraic varieties with* G_m *action*, Astérisque, **20** (1974) 1-131.

[W] K. Weierstrass, *Zur Theorie der Abelschen Functionen*, J. Reine Angew. Math., **47** (1854) 289-306.

Shigeki Matsutani:
8-21-1 Higashi-Linkan Minami-ku,
Sagamihara 252-0311,
JAPAN.
e-mail: rxb01142@nifty.com